

**GOLISANO
CHILDREN'S
HOSPITAL
AT STRONG**



**Pediatric Pain
Reference Cards**

*Pediatric Pain Subcommittee
University of Rochester*

Copyright 2005 The University of Rochester. These materials may not be reproduced without the expressed written permission of the University of Rochester.

SMH 1398 (Rev. 11/05)

Pediatric Pain Committee Members

Co-Chairs – Elise van der Jagt, MD, MPH
Kathy Rideout, PNP

Colleen Barron, CCLS, Child Life

Carol Buzzard, MD - Pediatric Cardiology

Keith DeMonte, PharmD - Clinical Pediatric Pharmacist

Irene Dutko Fioravanti, PNP - APN, 4-3600

Millee Francz, RN - AC6 Pain Resource Nurse

Michelle Gullace, PNP – APN, 4-1400/Palliative Care

Pam Hensel, PNP – APN, 4-2800

David Korones, MD - Pediatric Palliative Care

Pat Lamarche, PNP - APN, NICU

Marilyn Marano-Hyman, RN - NICU Pain Resource Nurse

Marcy Noble, RN - Pediatric Emergency Department

Julie Pietraszewski, PNP - Pediatric Surgery

Amy Roth, RN - 4-3600 Pain Resource Nurse

Marika Stone, MD - Anesthesia Pain Service

Amy Vallee, RN - 4-1400 Pain Resource Nurse

Tara Wengert, RN - 4-1600 Pain Resource Nurse

Golisano Children's Hospital at Strong Pediatric Acute Pain Algorithm

(See Pain Manual for NICU and Newborn Pain Protocols)

ASSESSMENT

1. Previous pain management including nonpharmacologic and pharmacologic interventions
2. Current pain complaint: Onset, location, duration, intensity, temporal pattern, aggravating factors, alleviating factors
3. Current pain medications/other medications
4. Contributing factors, e.g. course of disease, anxiety/fears, development, temperament, age
5. Risk factors, e.g. airway stability, disease process
6. Allergies/Sensitivities

**Assess pain using developmentally appropriate pain scale
and obtain pain rating (0 - 10)**

NONPHARMACOLOGIC OPTIONS

- **Physical Strategies:** massage; positioning, application of heat or cold; swaddling; sucrose pacifier; reduction of stimuli (noise control, dim lights, group care to decrease # of interactions).
- **Cognitive Strategies:** reassurance; distract by using art, play, child life activities, and music.
- **Child Life Specialist** for consultation to assist with coping strategies and/or diversional activities.
- **Psychological Evaluation** to assess if patient is a candidate for psychological interventions and/or possible self-regulatory strategies.

See Adjunctive Pain Management Strategies in Pain Manual

PHARMACOLOGIC OPTIONS (See Pain Manual for further dosing recommendations)

MILD PAIN (1-3)	MODERATE PAIN (4-7)	SEVERE PAIN (8-10)
<ul style="list-style-type: none">• Acetaminophen PO/PR• Ibuprofen PO• Morphine (Starting dose: 0.025 mg/kg/dose) IV (for patients unable to have PO/PR meds)	<ul style="list-style-type: none">• Ketorolac PO/ IV (>1 year/age) - d/c oral NSAIDS• Codeine PO• Oxycodone PO• Hydrocodone/ Acetaminophen PO (d/c Acetaminophen)• Morphine (PO - immediate release)• Hydromorphone PO• Morphine IV (Starting dose: 0.05 mg/kg/dose or 0.02 mg/kg/hour)• Morphine PCA (if age appropriate)• Hydromorphone IV	<ul style="list-style-type: none">• Morphine IV (starting doses)<ul style="list-style-type: none">• interval dosing (0.1 mg/kg/dose)• continuous dosing (0.05 - 0.1 mg/kg/hr)• Morphine PCA (add basal rate or increase total mg/hr)• Hydromorphone IV or PCA• Fentanyl IV or PCA (consider Anesthesia Pain Service Consult)• Anesthesia Pain Service Consult

Add NSAIDS and/or Acetaminophen, round the clock, in combination with any of the above medications, if not contraindicated

Reassess at appropriate intervals

GUIDELINES

- 1. Use Pain/Sedation Resource Manual for all analgesic/sedative dosing.**
- Evaluate pain on all patients using a 0 - 10 scale
 - A. Mild pain: 1 - 3
 - B. Moderate pain: 4 - 7
 - C. Severe pain: 8 - 10
- For chronic moderate or severe pain:
 - A. Give baseline medication around the clock
 - B. For breakthrough pain:
 - Continuous IV infusion: start at 50% of hourly dose and administer q 30-60 minutes. Dosing q 15 minutes may be necessary for some patients.
 - Intermittent IV dosing: 10% of total daily dose q 30-60 minutes.
 - Oral: 10% of total daily dose as a PRN given q 1-2 hours.
 - C. Adjust baseline upward daily in amount roughly equivalent to total amount of PRN
 - D. Negotiate with patient target level of relief, but usually at least achieving level < 4.
- In general, oral route is preferable, then transcutaneous > subcutaneous > intravenous.
- When converting from one opioid to another, some experts recommend reducing the equianalgesic dose by 1/3 to 1/2, then titrate as in #3 above.
- Infants < 6 months or those with severe renal or liver disease, should start on 1/4 to 1/2 the usual starting dose.
- If parenteral medication is needed for mild pain, use half the usual starting dose of morphine or equivalent.
- Naloxone (Narcan) should only be used in emergencies:
 - Dilute naloxone (0.4 mg/ml) 0.1 mg (0.25 ml) with 9.75 ml NS (total volume 10 ml)
 - Give 5.0 mcg/kg (0.5 ml) slow IVP until effect
 - Monitor patient q 15 minutes
 - May need to repeat again in 30-60 minutes
- Short-acting preparations should be used acutely & post-op. Switch to long-acting preparations when pain is chronic and the total daily dose is determined.
- For children < 1 yr (approx 10 kg), place on AB monitor when receiving parenteral narcotics (also consider for children with developmental disabilities, h/o prematurity and known respiratory difficulties).

Information adapted from *Facts and Comparisons 1997* and *APS Principles of Analgesic Use in the Treatment of Acute Pain and Cancer Pain* (4th Ed.) 1999.

Pediatric Pain Rating Scales

Premature Infant Pain Profile (PIPP) *(Stevens, et al, 1996)*

	Process	Indicator	0	1	2	3
Modifying Factors	Chart	Gestational Age (at time of observation)	36 wks and more	32 wks to 35 6/7 wks	28 wks to 31 6/7 wks	Less than 28 wks
	Observe Infant 15 sec Observe baseline Heart Rate Oxygen Saturation	Behavioral State	Active/ Awake Eyes open Facial movements	Quiet/ Awake Eyes open No facial movements	Active/ Sleep Eyes Closed Facial movements	Quiet/ Sleep Eyes Closed No facial movements
Behavioral/Physiologic Factors	Observe Infant 30 sec	Heart Rate Max	0-4 beats/min increase	5-14 beats/min increase	15-24 beats/min increase	25 beats/min or more increase
		Oxygen Sat Min	1.9% decrease	2-4% decrease	5-7% decrease	Greater than 8% decrease
		Brow bulge	None	Minimum	Moderate	Maximum
		Eye Squeeze	None	Minimum	Moderate	Maximum
		Nasolabial Furrow	None	Minimum	Moderate	Maximum

PIPP Pain Assessment Score and Treatment

Score	0-6	7-12	13-21
Therapy	Employ non-pharmacologic measures	Consider non-narcotic analgesia and/or Employ non-pharmacologic measures	Treat with narcotic analgesia and/or Employ non-pharmacologic measures

FLACC Scale *(Merkel, et al, 1997)*

Children < 3 years and Children with Developmental Disabilities

Categories	0	1	2
Face	No particular expression or smile	Occasional grimace or frown, withdrawn, disinterested	Frequent to constant quivering chin, clenched jaws
Legs	Normal position or relaxed	Uneasy, restless, tense	Kicking, or legs drawn up
Activity	Lying quietly, normal position, moves easily	Squirming, shifting back and forth, tense	Arched, rigid or jerking
Cry	No cry (awake or asleep)	Moans or whimpers; occasional complaint	Crying steadily, screams or sobs, frequent complaints
Consolability	Content, relaxed	Reassured by occasional touching, hugging or being talked to, distractible	Difficult to console or comfort

- Each of the five categories is scored from 0 - 2, which results in a total score between 0 - 10.
- Document the total score by adding numbers from each of the five categories.

Wong-Baker FACES Pain Rating Scale *(Wong, et al, 1999)*

Children > 3 years



0

No Hurt



2

Hurts Little Bit



4

Hurts Little More



6

Hurts Even More



8

Hurts Whole Lot




10

Hurts Worst

Numeric Scale

Teenagers and Young Adults



MEDICATION	EQUIANALGESIC DOSE (for chronic dosing)	
	IM/IV onset 15-30 min	PO onset 30-60 min
MORPHINE	10 mg	30 mg
OXYCODONE	Not Available	20 mg
HYDROMORPHONE (Dilaudid)	1.5 mg	7.5 mg
METHADONE	10 mg	24 hr Oral Morphine Oral Morphine: <u>Morphine</u> <u>Methadone Ratio</u> <30 mg 2:1 31-99 mg 4:1 100-299 mg 8:1 300-499 mg 12:1 500-999 mg 15:1 > 1000 mg 20:1
FENTANYL	100 mcg (single dose) 200 mcg (cont infusion)	24 hr oral MS dose <u>Initial Patch</u> <u>MS dose</u> 25 mcg/hr 90 mg 50 mcg/hr 180 mg 100 mcg/hr 360 mg
MEPERIDINE (Demerol)	75 -100 mg	300 mg
CODEINE (Tylenol #3) (Tylenol #4)	120 mg (IM only)	200 mg
HYDROCODONE (Vicodin, Lortab)	Not Available	30 mg

MED	USUAL STARTING DOSES Pediatric patients < 40 kg		COMMENTS (Not all dosage forms are available for inpatients. Consult unit pharmacist or CIS for availability)
	PARENTERAL	PO	
MORPHINE	Infant = 0.05-0.1 mg/kg/dose Child = 0.1-0.2 mg/kg/dose (Both given q 2-4 hours)	IR = 0.2-0.5 mg/kg/dose q 4-6 hrs CR = 0.3-0.6 mg/kg/dose q 8-12 hrs	Oral sol. (2 mg/ml): Conc. (20 mg/ml) can be given buccally. MSIR (Morphine Immediate-Release tablets - 15, 30 mg) Morphine sustained-release (15, 30, 60, 100, 200 mg) q 12 hrs. Use cautiously in severe renal disease.
OXY-CODONE	Not Available	0.05-0.15 mg/kg/dose q 4-6 hrs	Oxy IR (Oxycodone Immediate-Release tablets - 5 mg). Oxycodone sustained-release (10, 20, 40, 80 mg) q 12 hrs. Percocet (oxycodone/acetaminophen - Schedule III); 2.5/325, 5/325, 7.5/500, 10/650 mg). Monitor total acetaminophen dose. (For children <40 kg, 15 mg/kg/dose with maximum 1000 mg/dose; for >40 kg, maximum 1000 mg/dose and 4000 mg/day).
HYDRO-MORPHONE	0.015 mg/kg/dose q 3-6 hrs	0.03 - 0.08 mg/kg/dose q 3-6 hrs.	Tablets (2, 4, 8 mg) Oral liquid (5 mg/5 ml) Preferred for patients with renal disease
METHA-DONE	Not Available	Consult Pediatric Critical Care or Anesthesia Pain Service	Inexpensive. May help with myoclonus. Variable duration between individuals. Start doses q 6-12 hrs. and increase gradually. May accumulate with repetitive dosing (Days 2-5). Maximum dose 10 mg/dose. Conversion doses complex - consult Palliative Care or Anesthesia Pain Service.
FENTANYL	0.5-2 mcg/kg/dose q 30-60 minutes	25 mcg/hr q 72 hrs. (Transdermal) ⚡ Not recommended for opioid naïve)	IV: very short acting; associated with chest wall rigidity. Transdermal: See PDR for details of dose transition; include short-acting supplement for breakthrough pain; 12-hour delay onset and offset with patch.
MEPERIDINE	1-1.5 mg/kg/dose (Not to be used for analgesia; only use for shivering, procedure related pain)	Not Recommended	<u>Not recommended for standard analgesia.</u> May be used for shivering and procedural analgesia/sedation. Toxic metabolites accumulated with repeated doses, and with renal or hepatic disease. Contraindicated with MAOIs.
CODEINE	IM dosing not recommended IV dosing contraindicated	0.5-1.0 mg/kg/dose q 3-6 hrs.	Codeine alone - Schedule II prescription, all others Sched III. Tylenol #3 (codeine 30 mg w/ acetaminophen 300 mg). Tylenol #4 (codeine 60 mg w/ acetaminophen 300 mg). Tylenol w/codeine sol. (codeine 12 mg w/acet. 120 mg/5 ml). Monitor total acetaminophen dose. (For children <40 kg, 15 mg/kg/dose with maximum 1000 mg/dose; for > 40 kg, maximum 1000 mg/dose and 4000 mg/day).
HYDRO-CODONE	Not Available	0.2 mg/kg/dose q 4-6 hrs.	Vicodin (hydrocodone/acetaminophen: 5/500 mg). Vicoprofen (hydrocodone/bupropfen: 7.5 /200 mg). Lortab (hydrocodone/acet.: 2.5/500; 5/500; 7.5/500 mg). Norco (hydrocodone/acetaminophen: 10/325 mg). Monitor total acetaminophen dose. (For children <40 kg, 15 mg/kg/dose with maximum 1000 mg/dose; for > 40 kg, maximum 1000 mg/dose and 4000 mg/day).

M O D E R A T E T O S E V E R E

M I L D T O M O D E R A T E